

First case of nectarivory in *Podarcis liolepis* (Squamata: Lacertidae)

Grégory Deso^{1*}, Paul Simar Bayet¹, Paolo Varese², Ugo Schumpp³, and Marc Antoine Marchand⁴

Plant feeding in the genus *Podarcis* is known mainly from islands, where for example *Podarcis lilfordi* is known for its nectarivorous and frugivorous diet leading to pollination and seed dispersal (Pérez-Mellado and Casas, 1997; Pérez-Mellado et al., 2007). In France, a first case of nectarivory in the genus *Podarcis* has recently been reported on an islet in Marseille (Île du Grand Rouveau), with *P. muralis* (Laurenti, 1768) consuming nectar from *Lotus cytisoides* (Linné, 1753) (Cheylan and Rivière, 2020). Small islands can sometimes lack food resources, which forces reptiles to a high variability in their diet, ultimately using plant resources (Herrel et al., 2008). However, plant food interactions on the mainland are rarer, although some opportunistic *Podarcis* species may consume fruit from humans (Deso, 2014; Mačát et al., 2015). Recently, a case of florivory in *P. liolepis* (Boulenger, 1905) was reported in Spain where an individual pulled off and ingested a flower petal of *Lantana camara* (Linnaeus, 1753) (Van Den Berg, 2011). The Catalan lizard *Podarcis liolepis cebennensis* Guillaume & Geniez in Fretey, 1986 is a French endemic subspecies (Geniez and Crochet, 2003) widespread in the Occitan region but was discovered in 2007 in a new region (Provence-Alpes-Côte d'Azur), in the Vaucluse department, in the commune of Mornas, east of the Rhône river (Geniez et al., 2008). At present, only three populations of this species are known to exist

east of the Rhône River, which constitutes an originality and confers a high heritage value on these populations.

On 28 October 2022, a thermoregulating young male was observed at the foot of a mature pre-flowering shrub *Rhamnus alaternus* (Linné, 1753), during a targeted survey of the Mornas population which is found in cliff and scree habitats (WGS84: 44.205881°N, 4.730066°E). During the observation by two authors (GD and PSB), the young male went to the pre-flowering clumps of *Rhamnus alaternus* and carefully examined them (Fig. 1A) before lapping some of them (Fig. 1B). On closer inspection of the pre-flowering clumps, we realised that many buds were being pulled off and potentially consumed. The lizard spent about 25 seconds extracting juices from selected buds. It should be noted that, given the mild temperatures of autumn 2022, other plant species were also in pre-flowering or even flowering, such as *Juniperus oxycedrus* (Linné, 1753), *Vinca major* (Linné, 1753), *Lathyrus latifolius* (Linné, 1753) and *Saponaria oxymoides* (Linné, 1753). *Rhamnus alaternus* is a dioecious shrub typical of Mediterranean scrubland (Aronne and Wilcock, 1994). The male flowers produce a lot of pollen and the females a moderate amount of nectar (Ricciardelli d'Arbore and Intoppa, 2000). Its flowers and fruits are highly valued by hymenoptera, birds and mammals (Bas et al., 2006, 2009; Rico-Guzmán et al., 2012; Canale et al., 2015). This Mediterranean shrub, which is known for its antioxidant properties (Gadouche et al., 2022), provides a food resource for wildlife at a time of year when resources are scarce (Canale et al., 2015).

In order to identify the presence of sugary matter in pre-flower buds, we performed the flushing extraction protocol mentioned by Marrant et al. (2009); using a 0-90% HHTEC refractometer at an ambient temperature of 20 degrees. Nectar was sampled from 0.2 grams of pre-bud with a 2 ml wash of deionised water in a sterile 20 ml flask (protocol taken from Power et al., 2017). The flask was shaken for 1 minute. We replicated this protocol twice with two different groups of clumps in the autumn. According to Power et al. (2017), this

¹ Association Herpétologique de Provence Alpes Méditerranée, Maison des Associations, 384 route de Caderousse, 84100 Orange, France.

² Independent botanist, Luserna San Giovanni, 10062 Turin, Italy.

³ Conservatoire d'espaces naturels de Provence-Alpes-Côte d'Azur, 18 Avenue du Gand, 04200 Sisteron, France.

⁴ Conservatoire d'espaces naturels de Provence-Alpes-Côte d'Azur, Villa Thuret, 90 Chemin Raymond, 06160 Antibes, France.

* Corresponding author. E-mail: ahpam.contact@gmail.com



Figure 1. (A) Male *Podarcis liolepis* that is closely inspecting the pre-flowering bud of shrub *Rhamnus alaternus*. (B) Collection of nectar. Photographs by Grégory Deso.

technique is both effective and provides the most accurate data of fauna obtained resources when the floral clusters have a small amount of nectar. Results showed that the bud contents were positive for sugar presence, with a fairly low Brix measurement (1 degree Brix = 1 gr sucrose in Aronne and Malara, 2019) in autumn at 0.3 and 0.2% sugars in 2 ml of fluids. Using new DNA metabarcoding techniques (Martins et al., 2022), it would be interesting to carry out an analysis of faeces at several periods of the year in order to better understand the extent of the plant diet of *P. liolepis* on continental populations as suggested by Bassitta et al. (2022). As this lizard is particularly active throughout the year (Geniez and Cheylan, 2012) and the flowering of *Rhamnus alaternus* takes place partly in winter, it is possible that this Mediterranean shrub also provides it with a winter food resource, from the floral pre-buds. A better understanding of the role of fruit, nectar or other plant material in the diet of *P. liolepis* could help guide possible conservation actions.

Acknowledgements. We thank Philippe Geniez for his comments and Roger Meek for his pre peer review. We also acknowledge the comments and input of the anonymous reviewer of the journal.

References

- Aronne, G., Wilcock, C.C. (1994). Reproductive adaptability of *Rhamnus alaternus* L. (Rhamnaceae) in the Mediterranean environment. *Plant Biosystems* **128**: 67.
- Aronne, G., Malara, P. (2019): Fiber-optic Refractometer for In Vivo Sugar Concentration Measurements of Low-nectar-producing Flowers. *New Phytologist* **224**: 987–993.
- Bas, J.M., Pons, P., Gomez, C. (2006): Exclusive frugivory and seed dispersal of *Rhamnus alaternus* in the bird breeding season. *Plant Ecology* **183**: 77–89.
- Bas, J.M., Oliveras, J., Gómez, C. (2009): Myrmecochory and short-term seed fate in *Rhamnus alaternus*: Ant species and seed characteristics. *Acta Oecologica* **35**: 380–384.
- Bassitta, M., Alemany, I., Pérez-Mellado, V., Pérez-Cembranos, A., Navarro, P., Lluch, J., Jurado-Rivera, J.A., Castro, J.A., Picornell, A., Ramon, C. (2022): Phylogenetic, Microbiome, and Diet Characterisation of Wall Lizards in the Columbretes Archipelago (Spain): Clues for Their Conservation. *Diversity* **14**: 408. <https://doi.org/10.3390/d14050408>
- Canale, A., Benvenuti, S., Raspi, A., Benelli, G. (2015): Insect pollinators of the late winter flowering *Rhamnus alaternus* L., a candidate for honeybee-friendly scrubland spots in intensively managed agricultural areas. *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology: Official Journal of the Societa Botanica Italiana*.
- Cheylan, M., Rivière, V. (2020): Premier cas de nectarivorie chez *Podarcis muralis* (Laurenti, 1768) dans une population micro-insulaire française (île du Grand Rouveau, Var). *Bulletin de la Société Herpétologique de France* **174**: 101–103.
- Deso, G. (2014): Observation d'un comportement commensal

- anthropophile chez le Léopard sicilien des champs *Podarcis siculus campestris* (De Betta, 1857) en Toscane (Italie). Bulletin de la Société Herpétologique de France **153**: 57–60.
- Gadouche, L., Zerrouki, K., Zidane, A., Ababou, A., Bachir Elazaar, I., Merabet, D., Henniche, W., Ikhlef, S. (2022): Genoprotective, antimutagenic, and antioxidant effects of methanolic leaf extract of *Rhamnus alaternus* L. from the Bissa mountains in Algeria. Foods and Raw Materials **10**(2): 196–205.
- Geniez, P., Crochet, P.-A. (2003): Confirmation de l'existence, en France, de trois taxons méconnus: *Alytes obstetricans almogavarii* Arntzen & García Paris, 1995 (Amphibia, Discoglossidae), *Podarcis hispanica sebastiani* (Klemmer, 1964) (Reptilia, Lacertidae) et *Natrix natrix astreptophora* (Seoane, 1884) (Reptilia, Colubridae). Bulletin de la Société Herpétologique de France **105**: 41–53.
- Geniez, P., Deso, G., Jacob, L., Issartel, G. (2008): Thomas J.-P., Découverte de *Podarcis liolepis* (Boulenger, 1905) (Reptilia : Sauria : Lacertidae) en Provence, à l'est du Rhône (France). Bulletin de la Société Herpétologique de France **125**: 33–39.
- Geniez, P., Cheylan, M. (2012): Les Amphibiens et Reptiles du Languedoc Roussillon et Régions limitrophes. Atlas biogéographique, Biotope éditions, Mèze; Ouvrages scientifiques du Muséum national d'Histoire naturelle, Paris. (Collection Inventaires et biodiversité) 448 p.
- Herrel, A., Huyghe, K., Vanhooydonck, B., Backeljau, T., Breugelmans, K., Grbac, I., Van Damme, R., Irschick, D.J. (2008): Rapid large-scale evolutionary divergence in morphology and performance associated with exploitation of a different dietary resource. Proceedings of the National Academy of Sciences of the United States of America **105**: 4792–4795.
- Mačát, Z., Veselý, M., Jablonski, D. (2015): New case of fruit eating observation in *Podarcis siculus* (Rafinesque-Schmaltz, 1810) (Lacertidae) from Croatia. Biharean Biologist **9**: 158–159.
- Martins, B., Rocha, I.S., Rocha, R., Mata, V.A., Gonçalves, Y., Rato, C. (2022): Exotic supper: Exploring the diet composition of *Tarentola mauritanica* gecko in Madeira Island under a metabarcoding approach. Conference: 21st European Congress of Herpetology, Belgrade, Servia.
- Morant, D. S., Schumann, R., & Petit, S. (2009): Field methods for sampling and storing nectar from flowers with low nectar volumes. Annals of Botany **103**: 533–542.
- Pérez-Mellado, V., Casas, J.L. (1997): Pollination by a lizard on a Mediterranean island. Copeia **1997**: 593–595.
- Pérez-Mellado, V., Riera, N., Hernández-Estévez, J.A., Piccolo, V., Potter, C. (2007): A complex case of interaction between lizards and plants. The dead horse arum (*Dracunculus muscivorus*) and the Balearic lizard (*Podarcis lilfordi*). In: Corti, C., Lo Cascio, P., Biaggini, M., editors. Mainland and insular Lacertid Lizards: A Mediterranean Perspective. Firenze: Firenze University Press; 2007. p. 133–160.
- Ricciardelli d'Arbore, G., Intoppa, F. (2000): Fiori e api. Bologna: Calderini Edagricole 254 p.
- Rico-Guzmán, E., Terrones, B., Cantó, J.L., Bonet, A. (2012): Frugivore carnivores: preferences and contribution to seed dispersal of red fox *Vulpes vulpes* (Linnaeus, 1758) and stone marten *Martes foina* (Erxleben, 1777) in Carrascal de la Font Roja Natural Park (Alicante, Spain). Galemys **24**: 25–33.
- Van Den Berg, M. (2011): *Podarcis liolepis liolepis* (Boulenger, 1905) feeding on *Lantana camara* Linnaeus, 1753. Link: <https://www.lacerta.de/AS/Artikel.php?Article=124>.